

Toxicity of neem seed extract to *Tessaratomia papillosa* (Drury) relative to its allozyme genotypes

LU Fu-Ping, ZHAO Dong-Xiang*, LIU Ye-Ping, WANG Ai-Ping, CHEN Qing
(Environment and Plant Protection Institute, Chinese Academy of Tropical Agricultural Sciences, Danzhou, Hainan 571737, China)

Abstract: The relationships between the susceptibility to neem seed extract and the allozyme genotypes were examined in the first instar nymphs of *Tessaratomia papillosa* (Drury) for two polymorphic enzyme loci of *Pgi* and *Mdh* using allozyme analysis. Acute exposures of the insect to 5.2 mg/mL (LC_{50} value) neem seed extract resulted in 51.8% mortality in 24 h. Under the given experimental conditions, insect mortalities were significantly different among certain genotypes and alleles. At locus *Pgi*, the insects with the *Pgi-bb* genotype displayed the highest mortality (84%), whereas those with *Pgi-aa* and *Pgi-cc* showed the lowest mortalities (0 and 7%, respectively), which were significantly different from that of *Pgi-bb*. At locus *Mdh*, the insects with the genotype *Mdh-cc* and *Mdh-aa* exhibited the highest mortality (93%), but no mortality was observed in the insects with the genotype *Mdh-c*. These results clearly indicated that the insects with genotype *Mdh-aa* and *Mdh-cc* were significantly different from those with other three genotypes *Mdh-ab*, *Mdh-bb* and *Mdh-bc* in response to neem seed extract. In contrast, the mortalities of the insects with the *Pgi-a* and *Mdh-c* allele were the lowest, and were significantly different from those with other alleles. Our studies showed that individuals of *T. papillosa* with different genotypes had significantly different responses to neem seed extract. Such distinct relationships between the insect susceptibility to neem seed extract and its allozyme genotypes may allow us to use certain genotypes and alleles as genetic markers to assess the susceptibility of *T. papillosa* to neem seed extract.

Key words: *Tessaratomia papillosa*; neem seed extract; allozymes; genotype; allele; selective lethal effect

1 INTRODUCTION

Tessaratomia papillosa (Drury) is an important pest of lichi and longan. It mainly damages litchi, longan, and citrus fruit trees, and may cause great economic loss by decreasing the output and quality of the fruits if not properly prevented and cured. The bug sucks juices of tender tips, spica and young fruits of the trees by stinging them with its needle-shaped mouthpart, and causes the shedding of flower and fruits (Xie *et al.*, 2004). Being frightened, it shoots foul-smelling liquid that can burn or darken the flowers, tender tips, tender leaves and young fruits, and even lead to downy mildew (Xu *et al.*, 1993). For a long time, controlling of *T. papillosa* mainly depends on using synthetic insecticides. Although beneficial, extensive use of these chemicals has created public concern about their effects on the environment and on human health. Consequently, intensive efforts are being made to find alternatives, especially insecticides of plant origin, which are readily biodegradable, and perceived to be environmentally safe and ecologically acceptable.

Neem extract is one kind of the best bio-pesticides

and is used widely to control insect pests. Neem-based insecticides containing azadirachtin have been reportedly used to control over 400 species of insects belonging to the orders Diptera, Hymenoptera, Coleoptera, Lepidoptera, Orthoptera and Hemiptera, including important agricultural pests, such as armyworms, leafminers, aphids, and whiteflies (Schmutterer, 1990; Uleichs *et al.*, 2001; Haseeb *et al.*, 2004; Terezinha *et al.*, 2004; Shoil *et al.*, 2005). Neem-based insecticides should be ideal alternatives to synthetic insecticides for controlling fruit tree, vegetable and tea insect pests. Since it is selective, neem presents a less negative impact on the ecosystems, and its association with biological control has been paid to more and more attention (Terezinha *et al.*, 2004). However to our knowledge, neem-based insecticides have not been used in the control of *T. papillosa*. Could it be used to control *T. papillosa*? How effective is it in controlling *T. papillosa*? With these questions in mind, we tested the toxicity of neem seed extract to *T. papillosa*.

Many literatures focused on the lethal relationship between environmental contamination and aquatic organisms via allozyme analysis, and evaluated the

基金项目: 中国热带农业科学院科技基金资助项目(Rky 0514)

作者简介: 卢芙蓉, 女, 1978年生, 甘肃金塔人, 硕士, 专业方向为昆虫群体遗传学及遗传毒理学, E-mail: fuping_36@163.com

* 通讯作者 Author for correspondences, E-mail: dongxiangzh@163.com

收稿日期 Received: 2005-08-09; 接受日期 Accepted: 2006-02-05

effects of contamination on a population's genetic structure (Guttman, 1994; Fore *et al.*, 1995; Duan *et al.*, 2000a, 2000b, 2001). The results indicated that the genetic distance analysis could be a more sensitive tool in demonstrating the overall genetic disturbance caused by environmental change and potential use of genetic distances in these organisms as a bioindicator for monitoring environmental changes (Duan *et al.*, 2000a, 2000b, 2001). It had been used to resolve questions such as the identification of resistant and sensitive genotypes, the relationship between genetic diversity and tolerance, and how genotype and/or allele frequencies change after exposure to contaminants (Guttman, 1994). As an excellent reference, the theory and method had been adopted to study the correlation between insecticide toxicity and the allozyme genotypes and/or alleles of *Oxya chinensis* and *Plutella xylostella* (Li and Qiao, 2000; Li *et al.*, 2004a, 2004b, 2004c; Lu *et al.*, 2004a, 2004b). Researches have provided a new idea that the dynamic variability of allozyme genotypes and/or alleles can be taken as a potential genetic indicator to monitor pests' resistance to insecticides.

In this study, the toxicity of neem seed extract to the first instar nymphs of *T. papillosa* was tested, so that effects of neem-based insecticides on *T. papillosa* can be preliminarily evaluated. The allozyme analysis was used to assess the resistant risk of neem-based insecticides to *T. papillosa* by determining whether the insect's survival was associated with specific genotypes at selected enzyme loci during exposure. Malate dehydrogenase (MDH, E.C. 1.1.1.37) and glucose-6-phosphate isomerase (PGI, E.C. 5.3.1.9) were chosen because of their polymorphism and high activity in *T. papillosa* under our experiment conditions, and were extensively studied in allozyme analysis (Li *et al.*, 2004a, 2004b, 2004c; Lu *et al.*, 2004a, 2004b).

2 MATERIALS AND METHODS

2.1 Sample collection

The first instar nymphs of *T. papillosa* were collected from the research base of Chinese Academy of Tropical Agricultural Sciences in Danzhou, Hainan, and fed with fresh litchi leaves in insect net at room temperature before conducting acute toxicity treatment with neem seed extract.

2.2 Insecticide bioassays

The neem seed extract was provided by Laboratory of Insect Toxicology, South China Agricultural University.

Susceptibility of the first instar nymphs of *T. papillosa* to neem seed extract was evaluated using topical application. Five different concentrations (0.001, 0.003, 0.007, 0.01 and 0.03 g/mL) of

neem seed extract were prepared with ethanol as the solvent. The bioassay was carried out with five neem seed extract doses as treatments and a solvent as the control. Each treatment or control was repeated three times, and there were twenty individuals in each replicate. The insects were then put in Petri dishes with fresh litchi leaves at room temperature. Mortality was determined after 24 h.

2.3 Acute toxicity treatment

With the same topical application, a sample of first instar nymphs of *T. papillosa* was treated with the concentration of LC₅₀ neem seed extract based on the result of 2.2. After 24 h, dead and surviving individuals were stored at -80°C individually before electrophoresis.

2.4 Allozyme electrophoresis

Allozyme analysis was performed using horizontal starch gel electrophoresis (Richardsom, 1986; Wang, 1998; Li *et al.*, 2004). The 12.0% (w/v) gel was prepared using a mixture of soluble starch and potato starch (Sigma) at a ratio of 1:1 for electrophoresis. Phosphate buffer (0.05 mol/L, pH 8.0) was used as electrode buffer, and the ratio of electrode buffer to gel buffer was 9:1.

The leg muscle of each insect was removed and homogenized in 20 µL double distilled water on an ice pan.

The sample loading and gel staining were carried according to Murphy *et al.* (1996) and Wang (1998). Gels were run at constant voltage of 290 V for about 4 and a half hours with crushed ice as coolant at the top of gel at 4°C in a refrigerator.

Two polymorphic enzymes were examined: malate dehydrogenase (MDH, E.C. 1.1.1.37) and glucose-6-phosphate isomerase (PGI, E.C. 5.3.1.9). Alleles were identified by labeling the fastest migrating allele with *a*, the second fastest allele with *b*, and so on.

2.5 Data analysis

Concentration-mortality regression lines were maintained with SAS. The differential mortality among the allozyme genotypes and alleles of treated *T. papillosa* were compared by contingency table χ^2 test, herein, each genotype as a group and the individual number as replicate.

The genetic structures of the two allozyme genotypes were determined via BIOSYS-2 (Swofford *et al.*, 1981), with which allele frequency, percent polymorphic loci, heterozygosity (*H*), fixation index (*F*), goodness-of-fit to Hardy-Weinberg's (H-W) equilibrium and Roger's genetic distance (*D*) were calculated.

3 RESULTS

3.1 The toxicity determination

The mortality of the first instar nymphs of *T.*

papillosa exposed to neem seed extract depended upon the concentration of extract and a linear relationship was determind. The linear relationship was $y = 1.54 + 0.456x$, LC_{50} was 5.2 mg/mL and R^2 was 0.7804. The mortality of the first instar nymphs of *T. papillosa* at the lowest and the highest concentration was 16.67% and 88.33% , respectively.

3.2 The correlation analysis between mortality and genotypes

The mean mortality of 193 first instar nymphs of *T. papillosa* at the concentration of 5.2 mg/mL was 51.80% .

Overall , different lethal effect was observed at two different loci of different genotypes (Table 1). At the locus *Pgi* , significant differences were observed among the genotypes of high , medium and low mortality individuals. Individuals with *Pgi-bc* experienced the highest mortality(84%) ,but all individuals with *Pgi-aa* genotype survived. At the locus *Mdh* , *Mdh-aa* experienced the highest mortality , but all individuals with *Mdh-cc* genotype survived.

Table 1 Genotype effects on the probability of the two polymorphic loci *Pgi* and *Mdh* in *T. papillosa* population treated with neem seed extract

Locus	Mortality (%)				
	<i>aa</i>	<i>ab</i>	<i>bb</i>	<i>bc</i>	<i>cc</i>
<i>Pgi</i>	0 (29) a	58 (40) b	64 (61) b	84 (44) c	7 (15) a
<i>Mdh</i>	93 (15) a	54 (28) b	52 (96) b	61 (51) b	0 (13) c

Notes : The numbers in parenthesis were the total individual numbers including surviving and dead individuals with corresponding genotype. The data within a row followed with different letters were significantly different at 0.05 level. The same for Table 2.

3.3 The correlation analysis between mortality and alleles

The lethal effects of neem seed extract to *T. papillosa* were also different at the level of alleles (Table 2). χ^2 tests showed significant difference among the three alleles of *Pgi* . Allele *Pgi-b* that occurred in the most individuals exhibited the highest mortality ,

Pgi-a exhibited the lowest mortality , while the mortality of *Pgi-c* was medium. At *Mdh* locus , the mortalities of allele *Mdh-a* and *Mdh-b* showed no difference ($P < 0.05$) , but both showed a significant different lethal effect when compared with *Mdh-c* allele. However , the proportion of individuals with high mortality alleles was the highest of at least 75% at *Pgi* locus and 80% at *Mdh* locus.

Table 2 χ^2 tests for the mortality difference between *Pgi* and *Mdh* alleles of surviving and dead group of *T. papillosa* treated with neem seed extract

Locus	Mortality (%)		
	<i>a</i>	<i>b</i>	<i>c</i>
<i>Pgi</i>	23 (98) a	70 (226) b	52 (74) c
<i>Mdh</i>	61 (38) a	54 (271) a	40 (77) b

3.4 The allozyme genetic background analysis

When using the buffer system , the two enzymes all migrated from cathode to anode and each was found to have three allele polymorphic loci. At the two loci , the most frequent allele was the same , *Pgi-b* and *Mdh-b* (Table 3). The genotype frequencies at the two loci were all deviated significantly from Hardy-Weinberg equilibrium.

The frequency of the allele “ *b* ” had been elevated in the dead group in comparison with that of the initial group. The heterozygotes in dead groups were some exceeding since the *F* values in the two enzymes was lower than zero and the H_o was higher than H_e , but that was just opposite in alive and initial groups (Table 3 , 4).

The genetic identity (*I*) was higher between the alive and initial group as well as between the dead and initial group (Table 5) than that between the alive and dead group. The genetic distance (*D*) was greater between the alive and dead group than that between the alive and initial group as well as between the dead and initial group.

Table 3 Allele frequency and tests for Hardy-Weinberg (H-W) expectations at *Pgi* and *Mdh* loci of *T. papillosa* treated with neem seed extract

Alleles	<i>Pgi</i>			<i>Mdh</i>		
	Alive	Dead	Initial	Alive	Dead	Initial
Sample size	93	100	193	93	100	193
<i>a</i>	0.425	0.115	0.264	0.081	0.115	0.098
<i>b</i>	0.366	0.690	0.534	0.672	0.730	0.702
<i>c</i>	0.210	0.195	0.202	0.247	0.155	0.199
χ^2	60.070**	16.073**	43.030**	19.025**	15.153**	17.212**
<i>F</i>	0.531	- 0.269	0.246	0.262	- 0.070	0.106

Initial : including survival and dead. ** $P < 0.01$; *F* : Fixation index.

Table 4 Genetic variability at the two loci in each *T. papillosa* group treated with neem seed extract

Groups	Mean sample size per locus	Mean number of alleles per locus	Percentage of loci polymorphic *	Mean heterozygosity	
				H_o	H_e^{**}
Alive	93.0 (0.0)	3.0 (0.0)	100	0.328 (0.027)	0.564 (0.081)
Dead	100.0 (0.0)	3.0 (0.0)	100	0.530 (0.070)	0.454 (0.022)
Initial	193.0 (0.0)	3.0 (0.0)	100	0.433 (0.023)	0.532 (0.074)

H_o : observed heterozygosity. H_e : Hardy-Weinberg expected heterozygosity. * A locus is considered polymorphic if the most common allele does not exceed 0.95 (standard errors in-parentheses); ** Unbiased estimate (see Nei , 1978).

Table 5 Nei (1978) unbiased genetic identity (*I*: below diagonal) and modified Roger’s genetic distance (*D*: above diagonal) among the three groups of *T. papillosa* treated with neem seed extract

Group	Alive	Dead	Initial
Alive	–	0.232	0.120
Dead	0.902	–	0.112
Initial	0.974	0.982	–

4 DISCUSSION

Data indicate that although the neem products contributed towards pest reduction , but lost its efficacy after treatment and pest population there after increased (Akbar *et al.* , 2003). It has been found that application of neem-based insecticide at recommended application rates do not harm aquatic invertebrates categorized as planktonic and filter feeding. The benthic invertebrate *Chironomus riparius* was , however , affected by multiple applications of neem. High concentrations of neem were possibly not economical once the resistance occurred (Awad , 2003). So , in the experiment , we also did resistance risking evaluation of *T. papillosa* to neem-based insecticide before it could be used to control the insect.

The results of the toxicity of neem seed extract suggested that the neem-based insecticide had efficacy on controlling *T. papillosa* , though such effect was not very well , because the linear slope was not sharp. Then we tested the tolerance of the insect to neem seed extract via correlation analysis between allozyme genotypes of *T. papillosa* and toxic effects of neem seed extract. We hope that an inspecting index and evaluating basis of *T. papillosa* resistance to neem-based insecticide can be provided.

The significant toxin-genotype/allele interaction suggested a differential tolerance of *T. papillosa* to neem seed extract among genotypes and alleles. *Pgi-bc*/*Pgi-aa* genotype was sensitive/tolerant to neem extract , and *Pgi-a* allele was tolerant to the toxicant. It was very interesting that such response of *T. papillosa* to neem seed extract was similar to that of *Oxya chinensis* to avermectin (Li *et al.* , 2004c). In addition to *Pgi* , *Mdh-aa* was sensitive to neem seed extract , but *Mdh-cc* and *Mdh-c* was the highest tolerant genotype and

alleles. It demonstrated the correlation between allozyme genotypes and/or alleles mortality of *T. papillosa* and toxic effects of neem seed extract. The neem seed extract had a selective lethal effect on the studied *T. papillosa* population (Table 1 , 2).

Insect resistance to pesticides is an evolutionary event , possibly forming the genetic structure alternations within a population under the constant directional selection by pesticides (Tang and Wu , 2000). Identifying the physiological mechanisms of resistance is the first step in the characterization of resistance genes and also a prerequisite for understanding the evolution of insecticide resistance and for resistance management (Tsagkarakou *et al.* , 2002). In the present study , the genotypes and alleles of the two enzyme PGI and MDH might not directly explain the resistance occurrence of *T. papillosa* to neem-based insecticides. But the correlation between the allozyme genotype and the toxic effects of neem seed extract had been demonstrated. Individuals with *Pgi-aa* , *Pgi-cc* and *Mdh-cc* genotypes had the highest tolerance to neem seed extract , and the alleles of *Pgi-a* and *Mdh-c* were the highest tolerant alleles to neem seed extract. It suggested that the increased these genotypes and alleles frequencies in the population will be useful as the potential resistant genetic marker of *T. papillosa* to neem-based insecticide.

In conclusion , to some extent , the neem seed extract had differentiation effect on the genetic structure of *T. papillosa* studied , but the effect was not significant as indicated by the high genetic identity and the low genetic distance between the alive and dead group (Table 5). Thus this may suggest that neem-based insecticides is not likely to cause resistance. The existence of homozygotic individuals in the *T. papillosa* population may be of benefit for them to survive the toxicity of neem-based insecticides. The increased frequency of the most frequent allele in the dead group in comparison with that of the alive and initial group may be a indirect indication for that the neem-based insecticide is an effective pesticide for control of *T. papillosa* .

Acknowledgement Funding for this study was provided by Chinese Academy of Tropical Agricultural Sciences grant Rky 0514 to ZHAO Dong-Xiang. We thank Prof. MA En-Bo(College of Life Science and Technology) , and Prof. DUAN Yi-Hao (College of Environmental Science and Resources , Shanxi University) for their ideas for allozyme

analysis of this study. We also thank HE Yan-Ping (Postdoctor of College of Agronomy and BioTechnology , China Agricultural University) for her manuscript edit during the revision process.

References

Akbar AR , Sarwar M , Moula B , Tofique M , 2003. Evaluation of synthetic and some plant origin insecticides against *Helicoverpa armigera* (Hübner) on chickpea. *Pakistan Journal of Biological Sciences* , 6(5) : 496 – 499 .

Awad OM , 2003. Operational use of neem oil as an alternative anopheline larvicide. Part B : environmental impact and toxicological potential. *Eastern Mediterranean Health Journal* , 9(4) : 646 – 658 .

Duan YH , Guttman SI , Oris JT , 2000a. Genotype and toxicity relationships among *Hyalella azteca* : I . Acute exposure to metals or low pH. *Environmental Toxicology and Chemistry* , 19(5) : 1 414 – 1 421 .

Duan YH , Guttman SI , Oris JT , Huang XD , Burton GA , 2000b. Genotype and toxicity relationships among *Hyalella azteca* : II . Acute exposure to fluoranthene-contaminated sediment. *Environmental Toxicology and Chemistry* , 19(5) : 1 422 – 1 426 .

Duan YH , Guttman SI , Oris JT , Bailer AJ , 2001. Differential survivorship among allozyme genotypes of *Hyalella azteca* exposed to cadmium , zinc or low pH. *Aquatic Toxicology* , 54 : 15 – 28 .

Fore SA , Guttman SI , Bailer AJ , Altferer DJ , Counts BV , 1995. Exploratory analysis of population genetic assessment as a water quality indicator. I . *Pimephales notatus* . *Ecotoxicol. Environ. Safety* , 30 : 24 – 35 .

Guttman SI , 1994. Population genetic structure and ecotoxicology. *Environ. Health Perspect.* , 102 : 97 – 100 .

Haseeb M , Liu TX , Jones WA , 2004. Effects of selected insecticides on *Cotesia plutellae* endoparasitoid of *Plutella xylostella* . *Bio. Control* , 49 : 33 – 46 .

Li CL , Duan YH , Lu FP , Guo YP , Ma EB , 2004a. Mortality differences among the *Oxya chinensis* genotypes at glucose-6-phosphate isomerase by trichlorphon. *Journal of Agro-Environment Science* , 23(2) : 381 – 383 . [李翠兰 , 段毅豪 , 卢芙萍 , 郭亚平 , 马恩波 , 2004a. 敌百虫对中华稻蝗磷酸葡萄糖异构酶基因型的致死性差异研究. 农业环境科学学报 , 23(2) : 381 – 383]

Li CL , Duan YH , Lu FP , Guo YP , Ma EB , 2004b. Lethal responses of allozyme genotypes of *Oxya chinensis* towards cyhalothrin. *Journal of Agro-Environment Science* , 23(3) : 444 – 447 . [李翠兰 , 段毅豪 , 卢芙萍 , 郭亚平 , 马恩波 , 2004b. 中华稻蝗等位酶基因型对农药氯氟氰菊酯致死性的响应. 农业环境科学学报 , 23(3) : 444 – 447]

Li CL , Duan YH , Lu FP , Guo YP , Ma EB , 2004c. Differential mortality among the allozyme genotypes of *Oxya chinensis* by pesticide avermectin. *Acta Genetica Sinica* , 31(11) : 1 241 – 1 247 . [李翠兰 , 段毅豪 , 卢芙萍 , 郭亚平 , 马恩波 , 2004c. 中华稻蝗等位酶基因型与阿维菌素急性死亡率差异研究. 遗传学报 , 2004 , 31(11) : 1 241 – 1 247]

Li CX , Ma EB , Zheng XY , 2004. Genetic structure of four geographic populations of *Locusta migratoria manilensis* in China. *Acta Entomologica Sinica* , 47(1) : 73 – 79 . [李春选 , 马恩波 , 郑先云 , 2004. 中国东亚飞蝗四个地理种群遗传结构的比较研究. 昆虫学报 , 47(1) : 73 – 79]

Li X , Qiao CL , 2000. Studies on the esterase allozyme of avamectin resistant diamondback moth (*Plutella xylostella* L.). In : Li DM ed. *Chinese Entomology towards the 21st Century*. Beijing : China Science Press 379 – 381 . [李暄 , 乔传令 . 2000. 抗阿维菌素小菜蛾酯酶等位酶的研究 . 见 : 李典谟主编 . 走向 21 世纪的中国昆虫学 . 北京 : 中国科学出版社 . 379 – 381]

Lu FP , Li CL , Duan YH , Guo YP , Ma EB , 2004a. The selective mortality of *Oxya chinensis* to malathion toxicity at allozyme genotypes. *Journal of Shanxi University* (Nat. Sci. Ed.) , 27(2) : 178 – 181 . [卢芙萍 , 李翠兰 , 段毅豪 , 郭亚平 , 马恩波 , 2004a. 马拉硫磷对中华稻蝗等位酶基因型的致死作用. 山西大学学报 , 27(2) : 178 – 181]

Lu FP , Li CL , Duan YH , Guo YP , Ma EB , 2004b. Impacts of malathion on population genetic structure of *Oxya chinensis* . *Hereditas (Beijing)* , 26(5) : 663 – 668 . [卢芙萍 , 李翠兰 , 段毅豪 , 郭亚平 , 马恩波 , 2004b. 马拉硫磷对中华稻蝗种群遗传结果的作用. 遗传 (北京) , 26(5) : 663 – 668]

Murphy RW , Sites JW , Buth DJ , Haufler CH , 1996. Protein I : isozyme electrophoresis. In : Hillis DM , Moritz C , Mable BK eds. *Molecular Systematics* (2nd ed.). Sunderland , MA , USA : Sinauer Associate , Inc. Publisher.

Nei M , 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* , 89 : 583 – 590 .

Richardsom BJ , 1986. Allozyme Electrophoresis : A Handbook for Animal Systematics and Population Studies. Academic Press. 145 – 215 .

Schmutterer H , 1990. Properties and potential of natural pesticides from the neem tree , *Azadirachta indica* . *Annual Review of Entomology* , 35 : 271 – 297 .

Shoil MG , Allant S , Liu TX , 2005. Effects of neem-based insecticides on beet armyworm (Lepidoptera : Noctuidae). *Insect Science* , 12 : 17 – 23 .

Swofford DL , Selander RB , 1981. BIOSYS-1 : a FORTRAN program for the comprehensive analysis of electrophoretic data in population genetics and systematics. *Journal of Heredity* , 72 : 419 – 426 .

Tang ZH , WU SX , 2000. Heredity and Evolution of Insect Resistance to Pesticides. Shanghai : Science and Technology Press. 283 – 323 . [唐振华 , 吴士雄 . 2000. 昆虫抗药性的遗传与进化. 上海 : 上海科学技术文献出版社 . 283 – 323]

Terezinha MDS , Nivania PC , Adalci LT , Júnior ALB , 2004. Effect of neem extract on the cotton aphid. *Pesq. Agropec. Bras.* , Brasília , 39(11) : 1 071 – 1 076 .

Tsagkarakou A , Pasteur N , Cuany A , Chevillon C , Navajas M , 2002. Mechanisms of resistance to organophosphates in *Tetranychus urticae* (Acari : Tetranychidae) from Greece. *Insect Biochemistry and Molecular Biology* , 32 : 417 – 424 .

Uleichs CH , Mewis I , Schnitzler WH , 2001. Efficacy of neem and diatomaceous earth against cowpea aphids and their deleterious effect on predating Coccinelidae. *Journal of Applied Entomology* , 125 : 571 – 575 .

Wang ZR , 1998. Plant Allozyme Analysis. Beijing : Science Press 74 – 163 . [王中仁 , 1998. 植物等位酶分析. 北京 : 科学出版社 . 74 – 163]

Xie QM , Liang GW , Zeng L , Lu YY , 2004. The life table of the experimental population of *Tessaratomia papillosa* on litchi tree caged. *Entomological Knowledge* , 40(1) : 34 – 35 . [谢钦铭 , 梁广文 , 曾玲 , 陆永跃 . 2004. 荔枝蜡的实验种群生命表. 昆虫知识 , 40(1) : 34 – 35]

Xu CF , Chen JY , Xia YH , Ke C , 1993. On transmission of longan witches ' broom by *Tessaratomia papillosa* Drury. *Acta Phytopathologica Sinica* , 24 : 284 . [许长藩 , 陈景耀 , 夏雨华 , 柯冲 , 1993. 荔枝蜡传播龙眼鬼帚病的研究. 植物病理学报 , 24 : 284]

印楝种子提取物对荔枝蝽的毒性及与其等位酶基因型之间的关系

卢芙萍, 赵冬香*, 刘业平, 王爱萍, 陈 青

(中国热带农业科学院环境与植物保护研究所, 海南儋州 571737)

摘要: 用 5.2 mg/ml(LC_{50}) 的印楝种子提取物对荔枝蝽 1 龄若虫进行急性毒性处理, 24 h 死亡率为 51.8%。通过等位酶分析检测了死亡与存活试虫两种酶(PGI 和 MDH), 两个基因座(*Pgi* 和 *Mdh*) 上各基因型及等位基因与印楝种子提取物毒性之间的关系, 进行致死性差异比较研究。结果表明, 印楝种子提取物对具有不同基因型及等位基因个体的致死性存在差异。在 *Pgi* 基因座上, *Pgi-bb* 基因型死亡率最高, 为 84%, *Pgi-aa* 和 *Pgi-cc* 基因型死亡率较低, 分别为 0 和 7%, 且与死亡率最高的 *Pgi-bb* 基因型存在显著差异($P < 0.05$)。在基因座 *Mdh* 上, *Mdh-aa* 基因型个体死亡率最高(93%), 而具有 *Mdh-cc* 基因型的个体全部存活了下来, 另外 3 个基因型 *Mdh-ab*、*Mdh-bb* 与 *Mdh-bc* 死亡率居中, 都与 *Mdh-aa*、*Mdh-cc* 基因型死亡率之间存在显著差异。在等位基因上, *Pgi-a* 和 *Mdh-c* 个体的死亡率都最低, 与各自其他两个等位基因的死亡率之间存在显著差异。结果说明不同基因型个体对印楝提取物具有不同的反应, 印楝种子提取物对荔枝蝽等位酶基因型及等位基因存在选择性致死作用。这种荔枝蝽对印楝种子提取物的敏感性与其等位酶基因型及等位基因之间显明的相关关系提示我们, 可将荔枝蝽种群中对印楝种子提取物敏感性低的基因型及等位基因作为遗传标记去监测荔枝蝽对印楝种子提取物的抗性状况。

关键词: 荔枝蝽; 印楝种子提取物; 等位酶; 基因型; 等位基因; 选择性致死作用

中图分类号: Q355 文献标识码: A 文章编号: 0454-6296(2006)02-0241-06

(责任编辑: 袁德成)